

Universal Banach spaces with a projectional resolution of the identity: category-theoretic approach

Wiesław Kubiś

Czech Academy of Sciences (Prague, Czech Republic)

and

Jan Kochanowski University (Kielce, Poland)

e-mail: wkubis@ujk.kielce.pl

Abstract

A classical result of Pełczyński [1] from 1969 says that there exists a separable Banach space P with a Schauder basis which is complementably universal for all separable Banach spaces with a basis, i.e. every space in this class is isomorphic to a complemented subspace of P . Two years later, Kadec [2] proved a similar result for Banach spaces with the *Banach approximation property* (BAP) – a sequence of projections onto finite-dimensional subspaces converging pointwise to the identity.

A straight generalization of a Schauder basis for a non-separable Banach space is the notion of a *Markushevich basis*, i.e. a bi-orthogonal system $\langle x_\alpha, y_\alpha \rangle_{\alpha \in \Gamma}$ such that $\{x_\alpha\}_{\alpha \in \Gamma}$ is linearly dense and $\{y_\alpha\}_{\alpha \in \Gamma}$ is total. It seems that for Banach spaces of the smallest uncountable density \aleph_1 , a natural requirement is that the space

$$\{y: y(x_\alpha) = 0 \text{ for all but countably many } \alpha\}$$

be norming. Such a basis is called *countably norming*. A Banach space X with a countably norming Markushevich basis has several nice properties. For instance: it admits a bounded 1-1 linear operator into a space of type $c_0(\Gamma)$ and it has an equivalent locally uniformly convex norm on X . The main tool for proving these properties is the notion of a *projectional resolution of the identity* (PRI), that is, a continuous transfinite sequence of pairwise compatible projections onto smaller subspaces, converging pointwise to the identity. An interesting fact is that, for a Banach space of the smallest uncountable density \aleph_1 , the existence of a PRI is actually equivalent to the existence of a countably norming Markushevich basis. Thus, in the class of spaces of density \aleph_1 , projectional resolution of the identity is a common generalization of BAP and Schauder basis.

We prove that, assuming the continuum hypothesis $2^{\aleph_0} = \aleph_1$, there exists a Banach space U of density \aleph_1 which has a projectional resolution of the identity and which is complementably universal for the class of all spaces with these properties. The space U has moreover some sort of homogeneity property for separable Banach spaces which makes it unique up to a linear isometry.

The space U is constructed by using purely category-theoretic methods. Namely, we only use the following two properties of the category $\mathfrak{B}_{\text{sep}}$ of separable Banach spaces with linear operators of norm ≤ 1 .

- (1) Every pair of left-invertible arrows in $\mathfrak{B}_{\text{sep}}$ with the same domain admits the pushout in $\mathfrak{B}_{\text{sep}}$.
- (2) Every countable inductive sequence consisting of left-invertible arrows in $\mathfrak{B}_{\text{sep}}$ has the colimit in $\mathfrak{B}_{\text{sep}}$.

The talk is based on the work [3].

References

- [1] A. PEŁCZYŃSKI, *Universal bases*, *Studia Math.* 32 (1969) 247–268
- [2] M.I. KADEC, *On complementably universal Banach spaces*, *Studia Math.* 40 (1971) 85–89
- [3] W. KUBIŚ, *Fraïssé sequences: category-theoretic approach to universal homogeneous objects*, preprint, <http://arxiv.org/abs/0711.1683>